

CLAIMS

What is claimed is:

5

1. A compound of Formula I



I

or a pharmaceutically acceptable salt thereof,

wherein:

10

Z is selected from:

HO₂C;

HO(H)N(O)C;

H(O)C-N(OH);

CH₃(O)C-N(OH);

15

CH₃(H)N(O)C-N(OH);

HS;











H₂N(O)₂S;


CH₃(H)N(O)₂S;

HO(O)P;

20

(HO)₂(O)P;





L is selected from:

- 5 C₃-C₅ alkylenyl;
- Substituted C₃-C₅ alkylenyl;
- 3- to 5-membered heteroalkylenyl; and
- Substituted 3- to 5-membered heteroalkylenyl;

10 Substituted L groups contain 1 or 2 substituents on a carbon atom or nitrogen atom independently selected from:

- HO;
- CN; and
- CF₃;

15 wherein each substituent on a carbon atom may further be independently F, and wherein 2 substituents may be taken together with a carbon atom to which they are both bonded to form the group C=O;

R¹ is independently selected from:

- C₅ or C₆ cycloalkylenyl-(C₁-C₈ alkylenyl);
- Substituted C₅ or C₆ cycloalkylenyl-(C₁-C₈ alkylenyl);
- 20 5- or 6-membered heterocycloalkylenyl-(C₁-C₈ alkylenyl);
- Substituted 5- or 6-membered heterocycloalkylenyl-(C₁-C₈ alkylenyl);
- Phenylenyl-(C₁-C₈ alkylenyl);
- Substituted phenylenyl-(C₁-C₈ alkylenyl);
- 5- or 6-membered heteroarylenyl-(C₁-C₈ alkylenyl);
- 25 Substituted 5- or 6-membered heteroarylenyl-(C₁-C₈ alkylenyl);
- Phenyl;

Substituted phenyl;
Naphthyl;
Substituted naphthyl;
5- or 6-membered heteroaryl;
5 Substituted 5- or 6-membered heteroaryl;
8- to 10-membered heterobiaryl; and
Substituted 8- to 10-membered heterobiaryl;

R^2 is independently selected from:

H;
10 C_1 - C_6 alkyl;
Phenyl- $(C_1$ - C_8 alkylene);
Substituted phenyl- $(C_1$ - C_8 alkylene);
Naphthyl- $(C_1$ - C_8 alkylene);
Substituted naphthyl- $(C_1$ - C_8 alkylene);
15 5- or 6-membered heteroaryl- $(C_1$ - C_8 alkylene);
Substituted 5- or 6-membered heteroaryl- $(C_1$ - C_8 alkylene);
8- to 10-membered heterobiaryl- $(C_1$ - C_8 alkylene);
Substituted 8- to 10-membered heterobiaryl- $(C_1$ - C_8 alkylene);
Phenyl-O- $(C_1$ - C_8 alkylene);
20 Substituted phenyl-O- $(C_1$ - C_8 alkylene);
Phenyl-S- $(C_1$ - C_8 alkylene);
Substituted phenyl-S- $(C_1$ - C_8 alkylene);
Phenyl-S(O)- $(C_1$ - C_8 alkylene);
Substituted phenyl-S(O)- $(C_1$ - C_8 alkylene);
25 Phenyl-S(O)₂- $(C_1$ - C_8 alkylene); and
Substituted phenyl-S(O)₂- $(C_1$ - C_8 alkylene);

Each substituted R^1 group contains from 1 to 3 substituents, and each substituted R^2 group contains from 1 to 4 substituents, wherein each substituent is independently on a carbon or nitrogen atom, independently selected from:

30 C_1 - C_6 alkyl;
CN;
CF₃;
HO;

(C₁-C₆ alkyl)-O;

(C₁-C₆ alkyl)-S(O)₂;

H₂N;

(C₁-C₆ alkyl)-N(H);

5

(C₁-C₆ alkyl)₂-N;

(C₁-C₆ alkyl)-C(O)O-(C₁-C₈ alkylenyl)_m;

(C₁-C₆ alkyl)-C(O)O-(1- to 8-membered heteroalkylenyl)_m;

(C₁-C₆ alkyl)-C(O)N(H)-(C₁-C₈ alkylenyl)_m;

(C₁-C₆ alkyl)-C(O)N(H)-(1- to 8-membered heteroalkylenyl)_m;

10

H₂NS(O)₂-(C₁-C₈ alkylenyl);

(C₁-C₆ alkyl)-N(H)S(O)₂-(C₁-C₈ alkylenyl)_m;

(C₁-C₆ alkyl)₂-NS(O)₂-(C₁-C₈ alkylenyl)_m;

3- to 6-membered heterocycloalkyl-(G)_m;

Substituted 3- to 6-membered heterocycloalkyl-(G)_m;

15

5- or 6-membered heteroaryl-(G)_m;

Substituted 5- or 6-membered heteroaryl-(G)_m;

(C₁-C₆ alkyl)-S(O)₂-N(H)-C(O)-(C₁-C₈ alkylenyl)_m; and

(C₁-C₆ alkyl)-C(O)-N(H)-S(O)₂-(C₁-C₈ alkylenyl)_m;

wherein each substituent on a carbon atom may further be independently selected from:


Halo; and


HO₂C;

wherein 2 substituents may be taken together with a carbon atom to which they are both bonded to form the group C=O;

25

wherein two adjacent, substantially sp² carbon atoms may be taken together with a diradical substituent to form a cyclic diradical selected from:





R is H or C₁-C₆ alkyl;

5 G is CH₂; O, S, S(O); or S(O)₂;

Each m is an integer of 0 or 1;

Q, when bonded to a nitrogen atom in group D, is selected from:

OC(O);

CH(R⁶)C(O);

10 OC(NR⁶);

CH(R⁶)C(NR⁶);

N(R⁶)C(O);

N(R⁶)C(S);



N(R⁶)C(NR⁶);

15 SC(O);


CH(R⁶)C(S);

SC(NR⁶);

C≡CCH₂;



20



Q, when bonded to a carbon atom in group D, is as defined above and may further be selected from:


- 5 OCH_2 ;
- $\text{N(R}^6\text{)CH}_2$;
- trans-(H)C=C(H) ;
- cis-(H)C=C(H) ;
- $\text{C}\equiv\text{C}$;
- $\text{CH}_2\text{C}\equiv\text{C}$;
- 10 $\text{CF}_2\text{C}\equiv\text{C}$;
- $\text{C}\equiv\text{CCF}_2$;










Each R^6 independently is H, $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_3\text{-C}_6$ cycloalkyl; 3- to 6-membered heterocycloalkyl; phenyl; benzyl; or 5- or 6-membered heteroaryl;


X is O, S, N(H), or N($\text{C}_1\text{-C}_6$ alkyl);

- 15 Each V is independently C(H) or N;




















D is a cyclic diradical group selected from:



































; and




























5



















5



















5












;




;




;




;




;




;




;



;




;




;


5




;




;




;




;



;



;



wherein the group D may be unsubstituted or substituted on a carbon atom or a nitrogen atom by replacement of a hydrogen atom with a group selected from:

CH₃;

CF₃;

C(O)H;

CN;

HO;

CH₃O;

C(F)H₂O;

C(H)F₂O; and

CF₃O;

wherein a carbon atom in the group D may further be substituted with F;

V¹ is a 5-membered heteroarylenyl containing carbon atoms and from 1 to 4

heteroatoms selected from 1 O, 1 S, 1 NH, 1 N(C₁-C₆ alkyl), and 4 N,

wherein the O and S atoms are not both present, and wherein the

5 heteroarylenyl may optionally be unsubstituted or substituted with 1
substituent selected from fluoro, methyl, hydroxy, trifluoromethyl, cyano,
and acetyl;

wherein each C₈-C₁₀ bicycloalkyl is a bicyclic carbocyclic ring that contains 8-, 9-

, or 10-member carbon atoms which are 5,5-fused, 6,5-fused, or 6,6-fused bicyclic

10 rings, respectively, and wherein the ring is saturated or optionally contains one
carbon-carbon double bond;

wherein each 8- to 10-membered heterobicycloalkyl is a bicyclic ring that

contains carbon atoms and from 1 to 4 heteroatoms independently selected from 2

O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4 N(H), and 4 N(C₁-C₆ alkyl), and wherein when two

15 O atoms or one O atom and one S atom are present, the two O atoms or one O
atom and one S atom are not bonded to each other, and wherein the ring is
saturated or optionally contains one carbon-carbon or carbon-nitrogen double
bond, and wherein the heterobicycloalkyl is a 5,5-fused, 6,5-fused, or 6,6-fused
bicyclic ring, respectively,

20 wherein each heterocycloalkyl is a ring that contains carbon atoms and from 1 to 4
heteroatoms independently selected from 2 O, 1 S, 1 S(O), 1 S(O)₂, 1 N, 4

N(H), and 4 N(C₁-C₆ alkyl), and wherein when two O atoms or one O

atom and one S atom are present, the two O atoms or one O atom and one

S atom are not bonded to each other, and wherein the ring is saturated or

25 optionally contains one carbon-carbon or carbon-nitrogen double bond;

wherein each heterocycloalkylenyl is a ring diradical that contains carbon atoms

and from 1 to 3 heteroatoms independently selected from 1 O, 1 S, 1 S(O),

1 S(O)₂, 1 N, 2 N(H), and 2 N(C₁-C₆ alkyl), and wherein when one O atom

and one S atom are present, the one O atom and one S atom are not bonded

30 to each other, and wherein the ring is saturated or optionally contains one
carbon-carbon or carbon-nitrogen double bond;

wherein each 5-membered heteroaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and each 6-membered heteroaryl contains carbon atoms and 1 or 2 heteroatoms independently selected from N, N(H), and N(C₁-C₆ alkyl), and 5- and 6-membered heteroaryl are monocyclic rings;

wherein a 5-membered heteroarylenyl is a 5-membered monocyclic diradical ring that contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, wherein the 1 O atom and 1 S atom are not both present, and 6-membered heteroarylenyl is a 6-membered monocyclic diradical ring that contains carbon atoms and 1 or 2 heteroatoms independently selected from 2 N;


wherein each heterobiaryl contains carbon atoms and from 1 to 4 heteroatoms independently selected from 1 O, 1 S, 1 N(H), 1 N(C₁-C₆ alkyl), and 4 N, and where the 8-, 9-, and 10-membered heterobiaryl are 5,5-fused, 6,5-fused, and 6,6-fused bicyclic rings, respectively, and wherein at least 1 of the 2 fused rings of a bicyclic ring is aromatic, and wherein when the O and S atoms both are present, the O and S atoms are not bonded to each other;

wherein with any (C₁-C₆ alkyl)₂-N group, the C₁-C₆ alkyl groups may be optionally taken together with the nitrogen atom to which they are attached to form a 5- or 6-membered heterocycloalkyl; and


wherein each group and each substituent recited above is independently selected.

2. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein Z is HO₂C.

3. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein Z is selected from:



4. The compound according to Claim 1, or a pharmaceutically acceptable salt thereof, wherein Z is selected from:



5. The compound according to any one of Claims 1 to 4, or a pharmaceutically acceptable salt thereof, wherein Q is $N(R^6)C(O)$.
6. The compound according to any one of Claims 1 to 4, or a pharmaceutically acceptable salt thereof, wherein Q is selected from:
- 10 $C\equiv C$;
 $CH_2C\equiv C$;
 $C\equiv CCH_2$;
 $CF_2C\equiv C$; and
 $C\equiv CCF_2$.
- 15 7. The compound according to Claim 1, selected from:
- 4-(3-{3-[3-(3,4-Difluoro-benzyl)-4-oxo-3,4-dihydro-quinazolin-6-yl]-prop-2-ynyl}-phenyl)-butyric acid; and
- 5-(3,4-Difluoro-benzyl)-7-methyl-4,6-dioxo-3a,4,5,6-tetrahydro-
- 20 thieno[3,2-c]pyridine-2-carboxylic acid [2-(3-mercapto-propoxy)-pyridin-4-ylmethyl]-amide;
- or a pharmaceutically acceptable salt thereof.
8. A pharmaceutical composition, comprising a compound according to
- 25 Claim 1, or a pharmaceutically acceptable salt thereof, admixed with a pharmaceutically acceptable carrier, excipient, or diluent.
9. The pharmaceutical composition according to Claim 8, comprising a compound according to Claim 7, or a pharmaceutically acceptable salt thereof,
- 30 admixed with a pharmaceutically acceptable carrier, excipient, or diluent.

10. A method for treating osteoarthritis, comprising administering to a patient suffering from osteoarthritis or rheumatoid arthritis a nontoxic effective amount of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof.

5

11. The method according to Claim 10, wherein the compound administered is a compound according to Claim 7, or a pharmaceutically acceptable salt thereof.

10